arXiv:0807.0155v1 [math.RT] 1 Jul 2008

On the unitarization of linear representations of primitive partially ordered sets

Roman Grushevoy and Kostyantyn Yusenko

Abstract. We describe all weights which are appropriated for the unitarization of linear representations of primitive partially ordered sets of finite type.

Mathematics Subject Classification (2000). Primary 65N30; Secondary 65N15.

Keywords. unitarizations, quivers, partially ordered sets, Dynkin graphs, subspaces, *-algebras, representations.

0. Introduction

The representation theory of partially ordered sets (posets) in linear vector spaces has became one of the classical field in linear algebra (see [1, 2, 3, 4] and other). See Section 1 for the details.

On other hand when one try to develop the similar theory for Hilbert spaces, one will be faced with difficulties even with three subspace two of which is orthogonal: it is impossible to obtain their description in reasonable way (it is so called $*$-wild problem) see [5, 6] . But adding linear relation

$$\alpha_1 P_1 + \ldots + \alpha_n P_n = \gamma I,$$

between the projections P_i on corresponding subspaces, in some cases gives nice answers which have deep interconnections with linear case (see Section 2 for the details).

The aim of this article is to show that each linear representation of primitive poset of finite type could be unitarized with some weight $\chi = (\alpha_1, \ldots, \alpha_n, \gamma)$ and to describe all possible weights χ appropriated to the unitarization for given linear representation of primitive poset (Section 3).

This work has been partially supported by the Scientific Program of National Academy of Sciences of Ukraine, Project No 0107U002333.

1. Posets and their linear representations

In this section we will briefly recall some results concerning to partially ordered sets and their linear representation.

1.1. Posets and Hasse diagrams

Let $(\mathcal{N}, \prec)$ be finite partially ordered set (or poset for short) which for us will be $\{1, \ldots, n\}$. By the width of the poset $\mathcal{N}$ we will understand the maximal number of two by two incomparable elements of this set.

The poset $\mathcal{N}$ of the width m called primitive and denoted by $(k_1, \ldots, k_m)$ if this set is the cardinal sum of m linearly ordered sets $\mathcal{N}_1, \ldots, \mathcal{N}_m$ with orders $k_1, \ldots, k_m$.

We will use the standard graphic representations for the poset $(\mathcal{N}, \prec)$ called Hasse diagram. This representation associate to each point $x \in \mathcal{N}$ the vertex v_x and the arrow $v_x \to v_y$ if $x \prec y$ and there is no $z \in \mathcal{N}$, such that $x \prec z \prec y$. For example let $\mathcal{N} = \{1, 2, 3, 4, 5, 6\}$ with the following order:

$$2 \prec 1, \quad 2 \prec 3, \quad 4 \prec 3, \quad 6 \prec 5,$$

then the corresponding Hasse diagram is the following:



For each primitive poset $\mathcal{N} = (k_1, ...k_m)$ its Hasse diagram has the following form:



1.2. Linear representations of posets. Indecomposability and Bricks

By the linear representation π of the poset $\mathcal{N}$ in some complex vector space V we will understand such correspondence to each element $i \in \mathcal{N}$ the subspace $V_i \subset V$, that $i \prec j$ implies $V_i \subseteq V_j$. We will denote the representation π in the following way: $\pi = (V; V_1, \ldots, V_n)$, where n is the cardinality of $\mathcal{N}$, and for the primitive posets $(k_1, \ldots, k_m)$ we will use the following notation $\pi = (V; V_1^{(1)}, \ldots, V_{k_1}^{(1)}; \ldots; V_1^{(m)}, \ldots, V_{k_m}^{(m)})$.

By the dimension vector d_π of the representation π we will understand the vector $d_\pi = (d_0; d_1, \ldots, d_n)$, where $d_0 = \dim V$, $d_i = \dim V_i$ (correspondingly for representations of primitive posets the dimensional vector will be denoted by $d_\pi = (d_0; d_1^{(1)}, \dots, d_{k_1}^{(1)}; \dots; d_1^{(m)}, \dots, d_{k_m}^{(m)}))$.

In fact the linear representations of poset $\mathcal{N}$ forms the additive category $\mathrm{Rep}(\mathcal{N})$, where the set of morphisms $\mathrm{Mor}(\pi_1, \pi_2)$ between two representations $\pi_1 = (V; V_1, \dots, V_n)$ and $\pi_2 = (W; W_1, \dots, W_n)$ consists of such linear maps $C : V \to W$, that $C(V_i) \subset W_i$. Two representations π_1 and π_2 of poset $\mathcal{N}$ are isomorphic (or equivalent) if there exists invertible morphism $C \in \mathrm{Mor}(\pi_1, \pi_2)$, i.e. there exist invertible linear map $C : V \to W$ such that $C(V_i) = W_i$.

One can define the direct sum $\pi = \pi_1 \oplus \pi_2$ of two objects $\pi_1, \pi_2 \in \mathcal{N}$ in the following way:

$$\pi = (V \oplus W; V_1 \oplus W_1, \dots, V_n \oplus W_n).$$

Using the notion of direct sum it is natural to define *indecomposable* representations as the representations that are not isomorphic to the direct sum of two non-zero representations, otherwise representations are called *decomposable.* It is easy to show that representation π is indecomposable iff there exist no non-trivial idempotents in $\mathrm{End}(\pi)$. The representation π is called *brick* if there is no non-trivial endomorphism of this representation. One can show that *brick* implies *indecomposability.* But there exist indecomposable representations of posets which are not 'brick': let us take

$$A_\alpha = \begin{pmatrix} 1 & \alpha \\ 0 & 1 \end{pmatrix}, \quad \alpha \in \mathbb{R} \backslash \{0\},$$

the corresponding representation $\pi_\alpha = (V; V_1, V_2, V_3, V_4)$ of the poset $\mathcal{N} = (1, 1, 1, 1)$

$$V = \mathbb{C}^2 \oplus \mathbb{C}^2; \quad V_1 = \mathbb{C}^2 \oplus 0, \quad V_2 = 0 \oplus \mathbb{C}^2,$$
$$V_3 = \left\{(x, x) \in \mathbb{C}^4 | x \in \mathbb{C}^2\right\}, \quad V_4 = \left\{(x, A_\alpha x) \in \mathbb{C}^4 | x \in \mathbb{C}^2\right\},$$

is indecomposable but is not brick.

We call the representation π *non-degenerated* if the components of dimension vector d_π satisfy the following conditions:

- $d_i \neq 0$;
- if $i \prec j$ then $d_i < d_j$;
- $d_i < d_0$,

otherwise the representation is called *degenerated.* Note that in works [3, 4] authors by term *non-degenerated* means just first and second from above conditions.

1.3. Posets and Quivers

To study the representations of the primitive posets it is helpful to use the well-developed representation theory of quivers (for the representations of quivers see for example [7, 8]).

A quiver $Q = (Q_0, Q_1, s, t : Q_1 \to Q_0)$ is given by a set of vertices Q_0, which for us will be $\{v_0, \dots, v_n\}$, and a set Q_1 of arrows. An arrow $\rho \in Q_1$ starts at the vertex $s(\rho)$ and terminates at $t(\rho)$. A representation X of Q is given by a vector space X_{v_i} for each vertex $v_i \in Q_0$ and linear operator $X_\rho : X_{s(\rho)} \to X_{t(\rho)}$. In similar way as for the posets one can define the set of morphisms between two representations X and Y, *indecomposable* and *bricks* representations (see [7] for the details).

To each poset $\mathcal{N}$ one can associate the quiver $Q^{\mathcal{N}}$ in the similar way as Hasse diagram was associated, the only difference is that we add the extra vertex v_0 to diagram and connect by arrows all vertices that corresponds to maximal elements of $\mathcal{N}$ with the vertex v_0. More formally, $Q_0^{\mathcal{N}} = \{v_0, v_x | x \in \mathcal{N}\}$ and $Q_1^{\mathcal{N}}$ consists of:

- the arrows $v_x \to v_y$ if $x \prec y$ and there are no such z that $x \prec z \prec y$, and
- the arrows $v_x \to v_0$ if there no such z that $x \prec z$.

For example for each primitive poset $\mathcal{N} = (k_1, \ldots k_m)$ the corresponding quiver $Q^{\mathcal{N}}$ has the following form:



Remark 1.1. Notice that the underlying graph of the quiver $Q_{\mathcal{N}}$ that corresponds to primitive poset $\mathcal{N} = (k_1, \ldots, k_m)$ is the tree which denoted by $T_{k_1+1,\ldots,k_m+1}$.

It is obvious that each linear representations π of the poset $\mathcal{N} = (k_1, \ldots, k_m)$ defines some representation X_π of the corresponding quiver $Q^{\mathcal{N}}$. Indeed, let be given the representation

$$\pi = (V; V_1^{(1)}, \ldots, V_{k_1}^{(1)}; \ldots; V_1^{(m)}, \ldots, V_{k_m}^{(m)})$$

of the poset $\mathcal{N}$ then the corresponding representation X_π of quiver $Q_{\mathcal{N}}$ could be built in the following way: to each vertex $v_i^{(j)}$ we associate the space $V_i^{(j)}$, $X_{v_i^{(j)}} = V_i^{(j)}$ (to the vertex v_0 we associate the space V, $X_{v_0} = V$) and to each arrow $v_i^{(j)} \to v_{i+1}^{(j)}$ we associate the embedding $V_i^{(j)} \hookrightarrow V_{i+1}^{(j)}$ (to the arrows $v_{k_i}^{(j)} \to v_0$ we associate the embedding $V_{k_i}^{(j)} \hookrightarrow V$). And vice versa every representation X of quiver Q with underlaying graph $T_{k_1+1,\ldots,k_m+1}$, such that for all $\rho \in Q_1$ X_ρ is monomorphism, defines a representation of the poset $\mathcal{N} = (k_1, \ldots, k_n)$:

$$\pi_X = (V; V_1^{(1)}, \ldots, V_{k_1}^{(1)}; \ldots; V_1^{(n)}, \ldots, V_{k_n}^{(n)}),$$

where $V = X_{v_0}$ it is a representation of the the vertex v_0, and $V_j^{(i)} = X_{v_j^{(i)} \to v_{j+1}^{(i)}} \cdot \ldots \cdot X_{v_{k_i}^{(i)} \to v_0}(X_{v_j^{(i)}})$ is an image of space corresponding to the vertex $v_j^{(i)}$ under the maps that corresponds to the path from vertex $v_j^{(i)}$ to vertex v_0.

In fact having take the representation of poset then using construction above build the representation of corresponding quiver and finally after building the representation of corresponding poset one will get the representation which is isomorphic to that we start on, i.e. $\pi \simeq \pi_{X_\pi}$ in the category $\mathrm{Rep}(\mathcal{N})$.

Proposition 1.2.

1. *The representations π_1 and π_2 of the poset $\mathcal{N}$ are isomorphic if and only if the corresponding representation X_{π_1} and X_{π_2} of the quiver $Q_{\mathcal{N}}$ are isomorphic.*
2. *The representation π is indecomposable if and only if the corresponding representation X_π is indecomposable.*

Proof.

1. The morphism $C : X_{\pi_1} \to X_{\pi_2}$, establish the isomorphism between the representations of the quiver $Q_{\mathcal{N}}$ if and only if the linear map C_{v_0} is a isomorphism between the representations π_1 and π_2 of $\mathcal{N}$.
2. If the representation π of primitive poset $\mathcal{N}$ is decomposable, i.e. $\pi \simeq \pi_1 \oplus \pi_2$, then the corresponding representation X_π of the quiver $Q = Q_{\mathcal{N}}$ has the following form: for all $v_i \in Q_0$ $X_{v_i} = U_i \oplus W_i$, for all $v_i \to v_j \in Q_1$ $X_{v_i \to v_j} = \hat{\Gamma}_{i,j} \oplus \tilde{\Gamma}_{i,j}$, where $\hat{\Gamma}_{i,j} : U_j \to U_i$ and $\tilde{\Gamma}_{i,j} : W_j \to W_i$, i.e. X is decomposable.

□

1.4. Finite type posets

Recall the the poset $\mathcal{N}$ has *finite (linear) representation type* if there exist only finitely many indecomposable representation of $\mathcal{N}$ in $\operatorname{Rep}(\mathcal{N})$. Result obtained by M.M.Kleiner [2] gives complete description of the posets of finite type. In case of primitive posets using connection between posets and quivers and analogical result for quivers (see [7], for instance) one can obtain description of the primitive posets of finite type.

Proposition 1.3. *Each primitive poset of finite type has one of the following form:*

- (k) *for all* $k \in \mathbb{N}$*;*
- (k_1, k_2) *for all* $k_1, k_2 \in \mathbb{N}$*;*
- $(k, 1, 1)$*, for all* $k \in \mathbb{N}$*;*
- $(2, 2, 1)$*;*
- $(3, 2, 1)$*;*
- $(4, 2, 1)$.

Proof. if

As it was shown in previous subsection that for every representation of primitive posets one can build the representation of corresponding quiver, non-isomorphic representations of the poset corresponds to non-isomorphic representations of the quiver. The quivers A_{k+1}, $A_{k_1+k_2+1}$, D_{k+3}, E_6, E_7, E_8 are the corresponding quiver for the posets listed in statement. Each of these quivers has finitely many of non-isomorphic indecomposable representations (see [7]), hence the listed posets also has finitely many non-isomorphic indecomposable representations.

only if One can show that each primitive poset that is not included in list contains subposet with corresponding extended Dynkin quiver. But this quivers has infinitely many non-isomorphic indecomposable representations (see [9, 10] and other for description). We list infinite series of non-isomorphic indecomposable representations of corresponding poset:

- for the poset $(1,1,1,1)$ (the corresponding quiver is $\tilde{D}_4$):

$$V=\mathbb{C}^2=\langle e_1,e_2\rangle; V_1=\langle e_1\rangle, V_2=\langle e_2\rangle, V_3=\langle e_1+e_2\rangle, V_4=\langle e_1+\lambda e_2\rangle, \lambda\neq 0,1;$$

- for the poset $(2,2,2)$ (the corresponding quiver is $\tilde{E}_6$):

$$V=\mathbb{C}^3=\langle e_1,e_2,e_3\rangle;$$
$$V_1=\langle e_1\rangle, V_2=\langle e_1,e_2\rangle; V_3=\langle e_3\rangle, V_4=\langle e_2,e_3\rangle;$$
$$V_5=\langle e_1+e_2+e_3\rangle, V_6=\langle e_1+e_2+e_3,\lambda e_1+e_2\rangle, \lambda\neq 0,1;$$

- for the poset $(3,3,2)$ (the corresponding quiver is $\tilde{E}_7$):

$$V=\mathbb{C}^4=\langle e_1,e_2,e_3,e_4\rangle;$$
$$V_1=\langle e_1\rangle, V_2=\langle e_1,e_2\rangle, V_3=\langle e_1,e_2,e_3\rangle;$$
$$V_4=\langle e_4\rangle, V_5=\langle e_3,e_4\rangle, V_6=\langle e_2,e_3,e_4\rangle;$$
$$V_7=\langle e_1+e_2+e_3,\lambda e_1+e_3+e_4\rangle, \lambda\neq 0,1;$$

- for the poset $(5,2,1)$ (the corresponding quiver is $\tilde{E}_7$):

$$V=\mathbb{C}^6=\langle e_1,e_2,e_3,e_4,e_5,e_6\rangle;$$
$$V_1=\langle e_1\rangle, V_2=\langle e_1,e_2\rangle, V_3=\langle e_1,e_2,e_3\rangle, V_4=\langle e_1,e_2,e_3,e_4\rangle, V_5=\langle e_1,e_2,e_3,e_4,e_5\rangle;$$
$$V_6=\langle e_5,e_6\rangle, V_7=\langle e_3,e_4,e_5,e_6\rangle;$$
$$V_8=\langle e_1+e_3+e_4+e_5,\lambda e_1+e_3+e_5+e_6,e_2+e_4\rangle, \lambda\neq 0,1;$$

here vectors e_i form basis in corresponding linear spaces.

Hence the theorem is proved. □

1.5. Coxeter functors for representations of posets in linear spaces

Coxeter functors for representations of quivers were studied in [12] for proving Gabriel's theorem.

Similar Coxeter functors F^+, F^- acting in the category $\operatorname{Rep}(\mathcal{N})$ of linear representations of posets was constructed in [4]. We are not going to give their full construction here, but we remind some basic necessary facts.

There are functors σ,ρ ($\sigma^2=\mathrm{id},\rho^2=\mathrm{id}$) which act between categories $\operatorname{Rep}(\mathcal{N})$ and $\operatorname{Rep}(\mathcal{N}^*)$ (where $\mathcal{N}^*$ is dual to poset $\mathcal{N}$). These functors change dimensions of representations by the following formulas:

$$(d_0;d_1^{(1)},\dots,d_{k_1}^{(1)};\dots;d_1^{(m)},\dots,d_{k_m}^{(m)})\xmapsto{\rho}(\sum_{j=1}^{m}d_{k_j}^{(j)}-d_0;d_{k_1}^{(1)},$$
$$d_{k_1}^{(1)}-d_1^{(1)},\dots,d_{k_1}^{(1)}-d_{k_1-1}^{(1)};\dots;d_{k_1}^{(m)},\dots,d_{k_m}^{(m)}-d_{k_m-1}^{(m)});$$

$$(d_0;d_1^{(1)},\dots,d_{k_1}^{(1)};\dots;d_1^{(m)},\dots,d_{k_m}^{(m)})\xmapsto{\sigma}(d_0;d_0-d_1^{(1)},\dots,d_0-d_{k_1}^{(1)};\dots;$$
$$d_0-d_1^{(m)},\dots,d_0-d_{k_m}^{(m)}).$$

By definition $F^+ = \rho\sigma,\ F^- = \sigma\rho$, hence their action on generalized dimensions could be written as follows:

$$(d_0; d_1^{(1)},\dots,d_{k_1}^{(1)};\dots;d_1^{(m)},\dots,d_{k_m}^{(m)}) \xmapsto{F^+} (\sum_{j=1}^{m} d_{k_j}^{(j)} - d_0; \sum_{j=2}^{m} d_{k_j}^{(j)} - d_0,$$

$$\sum_{j=2}^{m} d_{k_j}^{(j)} - d_0 + d_1^{(1)},\dots,\sum_{j=2}^{m} d_{k_j} - d_0 + d_{k_1-1}^{(1)};\dots;\sum_{j=1}^{m-1} d_{k_j}^{(j)} - d_0 + d_{k_m-1}^{(m)}); \tag{1.1}$$

$$(d_0; d_1^{(1)},\dots,d_{k_1}^{(1)};\dots;d_1^{(m)},\dots,d_{k_m}^{(m)}) \xmapsto{F^-} ((m-1)d_0 - \sum_{j=1}^{m} d_{k_j}^{(j)}; d_2^{(1)} - d_1^{(1)},$$

$$d_3^{(1)} - d_1^{(1)},\dots,d_{k_1-1}^{(1)} - d_1^{(1)}, d_0 - d_1^{(1)};\dots;d_{k_l-1}^{(m)} - d_1^{(m)}, d_0 - d_1^{(m)}). \tag{1.2}$$

Importance of these functors is due to the following fact (see [4]):

Any non-degenerate representation of finite type poset $\mathcal{N}$ is the following: $(F^+)^k\pi$, where π is degenerate.

2. Representations of posets in Hilbert spaces

In this section we will consider the unitary representation of the posets. We will show how these representations connected with $*$-representation of certain algebras that generated by projections and linear relations. The representations of these algebras associated with primitive posets could be studied using the Coxeter functors, which will allow us study unitary representations of the posets, using similar technique.

2.1. Unitary representations of posets

Denote by $\mathrm{Rep}(\mathcal{N},\mathrm{H})$ the sub-category in $\mathrm{Rep}(\mathcal{N})$. Its set of objects consists of finite-dimensional Hilbert spaces and two objects π and $\tilde{\pi}$ are equivalent in $\mathrm{Rep}(\mathcal{N})$ if there exists morphisms between them which is unitary operator $U : H \to \tilde{H}$ such that $U(H_i) = \tilde{H}_i$ (unitary equivalent). Representation $\pi \in \mathrm{Rep}(\mathcal{N},\mathrm{H})$ called *irreducible* iff C^*-algebra generated by set of orthogonal projections $\{P_i\}$ on the subspaces $\{H_i\}$ is irreducible. Let remark that indecomposability of representation π in $\mathrm{Rep}(\mathcal{N})$ implies irreducibility of $C^*(\{P_i, i \in \mathcal{N}\})$ but converse not true (see [11] for details).

The problem of classifying all irreducible objects in category $\mathrm{Rep}(\mathcal{N},\mathrm{H})$ becomes much harder even for primitive poset $\mathcal{N} = (1,2)$ it is impossible to classify in reasonable way all irreducible representations: indeed this lead us to classify up to unitary equivalence three subspace in Hilbert space, two of which are orthogonal, but it is well-known [5] that such task is $*$-wild. Hence it is natural to consider some additional relation.

Let us consider those objects $\pi \in \mathrm{Rep}(\mathcal{N}, \mathrm{H})$, $\pi = (H; H_1, \dots, H_2)$, which satisfy the following linear relation:

$$\alpha_1 P_1 + \ldots + \alpha_n P_n = \gamma I, \tag{2.1}$$

where α_i, γ are some positive real numbers. These objects form a category and we will also will denote it by $\mathrm{Rep}(\mathcal{N}, \mathrm{H})$ (this will not make problem, course in the sequel we will consider only those representations in Hilbert space of the posets, which satisfy (2.1)). If to fix the weight $\chi = (\alpha_1, \dots, \alpha_n; \gamma) \in \mathbb{R}_+^{n+1}$ and consider only representations satisfying (2.1)) with this fixed weight one obtain another category $\mathrm{Rep}_\chi(\mathcal{N}, \mathrm{H})$ which is subcategory in $\mathrm{Rep}(\mathcal{N}, \mathrm{H})$.

The surprising result of [13] is that in category $\mathrm{Rep}(\mathcal{N})$ each indecomposable representation equivalent to some object in $\mathrm{Rep}(\mathcal{N}, \mathrm{H})$ is brick.

2.2. Certain $*$-algebras and their representation in the category of Hilbert spaces

Let be given the weight $\chi = (\alpha_1, \dots, \alpha_n; \gamma) \in \mathbb{R}_+^{n+1}$, consider the following $*$-algebra $\mathcal{A}_{\mathcal{N},\chi}$, which is generated by n projections and corresponds to poset $\mathcal{N}$ of order n:

$$\begin{aligned} A_{\mathcal{N},\chi} := \mathbb{C}\langle p_1, p_2, \dots, p_n | p_i = p_i^2 = p_i^*, p_i p_j = p_j p_i = p_i, i \prec j, \\ \alpha_1 p_1 + \ldots + \alpha_n p_n = \gamma e \rangle. \end{aligned} \tag{2.2}$$

It's easy to see that every $*$-representation π of $*$-algebra $\mathcal{A}_{\mathcal{N},\chi}$ gives us representation of poset $\mathcal{N}$ in category $\mathrm{Rep}_\chi(\mathcal{N}, \mathrm{H})$ ($H_i = \pi(p_i)$), and vice versa every representation of poset $\mathcal{N}$ in category $\mathrm{Rep}_\chi(\mathcal{N}, \mathrm{H})$ generate the $*$-representation of $\mathcal{A}_{\mathcal{N},\chi}$. To put it another way, categories $\mathrm{Rep}_\chi(\mathcal{N}, \mathrm{H})$ and $\mathrm{Rep}(\mathcal{A}_{\mathcal{N},\chi})$ are equivalent.

For primitive poset $\mathcal{N} = (k_1, \dots, k_m)$ weights, for convenience, will be denoted by $\chi = (\alpha_1^{(1)}, \dots, \alpha_{k_1}^{(1)}; \dots; \alpha_1^{(m)}, \dots, \alpha_{k_m}^{(m)}; \gamma)$ and correspondingly

$$\begin{aligned} \mathcal{A}_{\mathcal{N},\chi} = \langle p_1^{(1)}, \dots, p_{k_1}^{(1)}, \dots, p_1^{(m)}, \dots, p_{k_m}^{(m)} | p_i^{(j)2} = p_i^{(j)*} = p_i^{(j)}, \\ p_i^{(j)} p_k^{(j)} = p_k^{(j)} p_i^{(j)} = p_i^{(j)}, \sum_{j=1}^{m} \sum_{i=1}^{k_l} \alpha_i^{(j)} p_i^{(j)} = \gamma e \rangle. \end{aligned} \tag{2.3}$$

Under some condition there exists isomorphisms between these algebras and $*$-algebras $\mathcal{A}_{\Gamma,\hat{\chi}}$ which are associated with star-shaped graphs $\Gamma = T_{k_1+1,\dots,k_m+1}$ and character $\hat{\chi} = (\beta_1^{(1)}, \dots, \beta_{k_1}^{(1)}; \dots; \beta_1^{(m)}, \dots, \beta_{k_m}^{(m)}; \gamma)$. $*$-Algebras $\mathcal{A}_{\Gamma,\hat{\chi}}$ defined in the following way:

$$\begin{aligned} \mathcal{A}_{\Gamma,\hat{\chi}} = \langle q_1^{(1)}, \dots, q_{k_1}^{(1)}, \dots, q_1^{(m)}, \dots, q_{k_m}^{(m)} | q_i^{(j)*} = q_i^{(j)}, q_i^{(j)} q_k^{(j)} = q_k^{(j)} q_i^{(j)} = \delta_{ik} q_i^{(j)}, \\ \sum_{j=1}^{l} \sum_{i=1}^{k_l} \beta_i^{(j)} q_i^{(j)} = \gamma e \rangle. \end{aligned} \tag{2.4}$$

Proposition 2.1. *Let $\mathcal{N}$ be primitive poset and $\chi = (\alpha_1^{(1)}, \dots, \alpha_{k_1}^{(1)}; \dots; \alpha_1^{(m)}, \dots, \alpha_{k_m}^{(m)}; \gamma)$ and $\hat{\chi} = (\beta_1^{(1)}, \dots, \beta_{k_1}^{(1)}; \dots; \beta_1^{(m)}, \dots, \beta_{k_m}^{(m)}; \gamma)$ be two given weights such that for all $j = 1, \dots, m$, $i = 1, \dots, k_j$ the following conditions hold:*

$$\beta_i^{(j)} = \sum_{s=i}^{k_j} \alpha_s^{(j)},$$

then there exist isomorphism ϕ between $*$-algebras $\mathcal{A}_{\mathcal{N},\chi}$ and $\mathcal{A}_{\Gamma,\hat{\chi}}$.

Proof. One can easily proof that map which is given on generators of algebras in the following way:

$$p_i^{(j)} \xmapsto{\phi} \sum_{s=1}^{i} q_s^{(j)}, \quad j = 1, \dots, l, i = 1, \dots, k_j,$$

establishes the isomorphism between $\mathcal{A}_{\mathcal{N},\chi}$ and $\mathcal{A}_{\Gamma,\hat{\chi}}$. □

The isomorphism ϕ can be extended to an equivalence functor between categories of representations of the algebras. Let π be representation of $*$-algebra $\mathcal{A}_{\Gamma,\hat{\chi}}$ then $\phi \circ \pi$ – representation of $\mathcal{A}_{\mathcal{N},\chi}$. Furthermore if π_1 and π_2 are equivalent objects in $\mathrm{Rep}(\mathcal{A}_{\Gamma,\hat{\chi}})$ then $\phi \circ \pi_1$ and $\phi \circ \pi_2$ are equivalent objects in $\mathrm{Rep}(\mathcal{A}_{\mathcal{N},\chi})$.

Let as recall some definitions from [15]:

Definition 2.2. An irreducible finite dimensional $*$-representation π of the algebra $\mathcal{A}_{\Gamma,\hat{\chi}}$ such that

$$\pi(q_i^{(j)}) \neq 0, j = 1, \dots, m, i = 1, \dots, k_j) \text{ and}$$
$$\sum_{i=1}^{k_j} \pi(q_i^{(j)}) \neq I, j = 1, \dots, m$$

will be called *non-degenerate*. By $\overline{\mathrm{Rep}}(\mathcal{A}_{\Gamma,\hat{\chi}})$ we will denote full subcategory of non-degenerate representations in the category $\mathrm{Rep}(\mathcal{A}_{\Gamma,\hat{\chi}})$.

Objects that corespondents to subcategory $\overline{\mathrm{Rep}}(\mathcal{A}_{\Gamma,\hat{\chi}})$ in $\mathrm{Rep}\mathcal{A}_{\mathcal{N},\chi})$ will be called non-degenerated as well. Non-degenerate representations of $*$-algebra $\mathcal{A}_{\mathcal{N},\chi}$ corresponds to non-degenerate representations of poset $\mathcal{N}$.

If one have degenerate representation of poset $\mathcal{N}$ in Hilbert space H, that is one of the following conditions are hold:

- There exist i such that $H_i = 0$ i.e. $P_i = 0$;
- There exist i such that $H_i = H_{i-1}$ i.e. $P_i = P_{i-1}$;
- There exist i such that $H_i = H$ i.e. $P_i = I$

In this case one can obtain non-degenerate representation of some subposet of $\mathcal{N}$ as follows. If for some i, $P_i = 0$ then one automatically have representation of poset $\mathcal{N}' = \mathcal{N} \setminus \{i\}$. If in representation π one have $P_i = P_{i-1}$ than one can consider π as representation of poset $\mathcal{N}'$ where $\pi(i-1) = H_i = H_{i-1}$ and $\sum_{j \in \mathcal{N} \setminus \{i-1,i\}} \alpha_j P_j + (\alpha_{i-1} + \alpha_i) P_{i-1} = \gamma I$. And finally, if for some i $\pi(i) =$

H one can consider representation π as representation of $\mathcal{N}'$ for which holds $\sum_{j\in\mathcal{N}'}\alpha_j P_j = (\gamma-\alpha_i)I$.

2.3. Coxeter functors for representations of $*$-algebras $\mathcal{A}_{\mathcal{N},\chi}$ in case of primitive poset $\mathcal{N}$

In work [14] there were developed Coxeter functors for representations of quivers in Hilbert spaces which prescribe so-called 'orthoscalarity' condition. It was proved that any irreducible representation of quiver which underlying graph is Dynkin diagram can be obtained from simplest ones by using this Coxeter functors.

These functors allows to construct some functors Φ^+ and Φ^-, called by Coxeter too, which act on representations of algebras $\mathcal{A}_{\Gamma,\hat{\chi}}$. Recall that $*$-algebras $\mathcal{A}_{\Gamma,\hat{\chi}}$ and their involutive representations are studied in many recent works (see [15, 17] and other). Using the isomorphism ϕ between $*$-algebras $\mathcal{A}_{\Gamma,\hat{\chi}}$ and $\mathcal{A}_{\mathcal{N},\chi}$ we obtain two new functors which we will be also denoted by Φ^+ and Φ^- and called Coxeter functors. These functors act from category $\mathrm{Rep}(\mathcal{A}_{\mathcal{N},\chi})$ to $\mathrm{Rep}(\mathcal{A}_{\mathcal{N},\chi_{\Phi^+}})$ and $\mathrm{Rep}(\mathcal{A}_{\mathcal{N},\chi_{\Phi^-}})$ respectively, where

$$\chi_{\Phi^+} = (\gamma-\sum_{i=1}^{k_1}\alpha_i^{(1)},\alpha_1^{(1)},\ldots,\alpha_{k_1-1}^{(1)};\ldots;$$

$$\gamma-\sum_{i=1}^{k_m}\alpha_i^{(m)},\alpha_1^{(m)},\ldots,\alpha_{k_m-1}^{(m)};(m-1)\gamma-\sum_{j=1}^{m}\alpha_{k_j}^{(j)});$$

$$\chi_{\Phi^-} = (\alpha_2^{(1)},\alpha_3^{(1)},\ldots,\alpha_{k_1}^{(1)},\sum_{j=2}^{m}\sum_{i=1}^{k_j}\alpha_i^{(j)}-\gamma;\ldots;$$

$$\alpha_2^{(k_m)},\ldots,\sum_{j=1}^{m-1}\sum_{i=1}^{k_j}\alpha_i^{(j)}-\gamma;\sum_{j=1}^{m}\sum_{i=1}^{k_j}\alpha_i^{(j)}-\gamma).$$

In other words, this functors can be considered as functors from $\mathrm{Rep}_\chi(\mathcal{N},\mathrm{H})$ to $\mathrm{Rep}_{\chi_{\Phi^\pm}}(\mathcal{N},\mathrm{H})$. Dimensions of representations are changed by formulas similar to (1.2) and (1.1) respectively.

3. Unitarization of linear representations of primitive posets

This section is devoted to the interconnection between linear and unitary representation of the primitive posets. Actually for primitive posets of finite type we will obtain the complete list of possible weights that appropriate to given indecomposable linear representation.

3.1. Unitarization

Let be given the representation $\pi \in \operatorname{Rep}(\mathcal{N})$, $\pi = (V; V_1, \ldots, V_n)$ of the poset $\mathcal{N}$. One say that π can be unitarized if there exists appropriate choice of hermitian structure $\langle\cdot,\cdot\rangle_{\mathbb{C}}$ in V, that corresponding projection $P_i : V \to V_i$ satisfy the following relation:

$$\alpha_1 P_1 + \ldots + \alpha_n P_n = \gamma I,$$

for some weight $\chi = (\alpha_1, \ldots, \alpha_n; \gamma) \in \mathbb{R}_+^{n+1}$, and correspondingly we say that π can be unitarized with weight χ if this weight is fixed.

Recall that the similar notion of *unitarization* was provided in article [14] for the unitarization of representations of given quiver Q. It fact that work studied the question when it is possible to define hermitian structure of each space $X_i, i \in Q_0$, in such way that in each vertex i the following condition satisfies:

$$\sum_{j \longrightarrow i} X_{j \longrightarrow i} X^*_{i \longrightarrow j} + \sum_{i \longrightarrow j} X^*_{i \longrightarrow i} X_{j \longrightarrow j} = \alpha_i I_{X_i},$$

where $X^*_{i \longrightarrow j}$ denotes the adjoint map to $X_{j \longrightarrow i}$ with respect to hermitian structure in X_i and X_j. One of the results of paper [14] is that if Q is a Dynkin quiver then every representation could be unitarizable, and if Q is an extended Dynkin quiver then there are representations that cannot be unitarized.

One can obtain analogical fact for representations of primitive poset of finite type using Coxeter functos. Noticed that it can be obtained using results from [14].

Proposition 3.1. *Every indecomposable linear representation of primitive poset of finite type can be unitarized with some weight.*

Proof. To start with, remark that it is obvious that any one dimensional representation with dimension vector $d_\pi = (1; d_1, \ldots, d_n)$ of any poset can be unitarized with weight $\chi = (\alpha_1, \ldots, \alpha_n; \gamma)$ satisfied *trace* condition $\sum_{s \in \mathcal{N}} d_s \alpha_s = \gamma$. All indecomposable representations of primitive posets $\mathcal{N} = (k), k \in \mathbb{N}$ and $\mathcal{N} = (k_1, k_2)$, $k_1, k_2 \in \mathbb{N}$ are one-dimensional hence they could be unitarized.

Let π be some representation of primitive poset $\mathcal{N}$ of finite type. There are two possibilities: π is degenerate or π is non-degenerate. In first case the representation π can be considered as one of some subposet $\mathcal{N}'$ of $\mathcal{N}$ than by induction it unitarize and corresponding unitary representation of $\mathcal{N}'$ can be restricted to unitary representation of $\mathcal{N}$ which is unitarization of π.

In the later case representation π by theorem from [4] which was mentioned in section 1.5 is as follows $\pi = (F^+)^k \pi'$ that is $(F^-)k\pi = \pi'$ where π' is degenerate. As degenerate representation π' unitarize with some weight χ'. Using functor $(\Phi^+)^k$ to corresponded unitary representation we obtain unitary representation with some weight χ equivalent to π. That is π unitarize with weight χ. As the result we obtain statement of the theorem and even more: this gives the algorithm which allows to describe all possible weights with which given representation of finite type primitive poset can be unitarized. □

Next theorem gives complete list of all possible weights for representations of primitive posets of finite type. In other words it describes the set of weights χ for every primitive poset $\mathcal{N}$ such that $*$-algebra $\mathcal{A}_{\mathcal{N},\chi}$ has indecomposable representation in fixed dimention D. Analogical result for $*$-algebras $\mathcal{A}_{\Gamma,\hat{\chi}}$ were obtained in [15] and following theorem can be obtained using result of [15] but we did it independently. The list of weights are organized in the following way: for each representation of primitive poset $\mathcal{N} = (k_1, \dots, k_n)$ (which for us is given by generalized dimensional $d = (d_1^{(1)}, \dots, d_{k_1}^{(1)}; \dots; d_1^{(n)}, \dots, d_{k_n}^{(n)}; d_0)$, since this gives a representation up to isomorphism) we state the condition on weights $\chi = (\alpha_1, \dots, \alpha_{k_1}; \beta_1, \dots; \gamma)$, under which it is possible to unitarize the linear representation.

Theorem 3.2. *For primitive poset $\mathcal{N}$ its linear representation with dimension D unitarize with every weight χ for which conditions C are satisfied:*

1. Poset $\mathcal{N} = (1;1;1)$



Dimensions D	*Conditions C*
$(0;0;0;1)$	$\gamma = 0$
$(0;0;1;1)$	$\delta = \gamma$
$(0;1;0;1)$	$\beta = \gamma$
$(0;1;1;1)$	$\beta + \delta = \gamma$
$(1;0;0;1)$	$\alpha = \gamma$
$(1;0;1;1)$	$\alpha + \delta = \gamma$
$(1;1;0;1)$	$\alpha + \beta = \gamma$
$(1;1;1;1)$	$\alpha + \beta + \delta = \gamma$
$(1;1;1;2)$	$\alpha < \gamma, \beta < \gamma, \delta < \gamma, \alpha + \beta + \delta = 2\gamma$

2. Poset $\mathcal{N} = (2;1;1)$



Dimensions	*Conditions*
$(0,0;0;0;1)$	$\gamma = 0$
$(0,1;0;0;1)$	$\alpha_2 = \gamma$
$(1,1;0;0;1)$	$\alpha_1 + \alpha_2 = \gamma$
$(0,0;1;0;1)$	$\beta = \gamma$
$(0,1;1;0;1)$	$\alpha_2 + \beta = \gamma$
$(1,1;1;0;1)$	$\alpha_1 + \alpha_2 + \beta = \gamma$
$(0,0;0;1;1)$	$\delta = \gamma$

$(0,1;0;1;1)$	$\alpha_2+\delta=\gamma$
$(1,1;0;1;1)$	$\alpha_1+\alpha_2+\delta=\gamma$
$(0,0;1;1;1)$	$\beta+\delta=\gamma$
$(0,1;1;1;1)$	$\alpha_2+\beta+\delta=\gamma$
$(1,1;1;1;1)$	$\alpha_1+\alpha_2+\beta+\delta=\gamma$
$(0,1;1;1;2)$	$\alpha_2<\gamma,\beta<\gamma,\delta<\gamma,\alpha_2+\beta+\delta=2\gamma$
$(1,1;1;1;2)$	$\alpha_1+\alpha_2<\gamma,\alpha_1+\alpha_2<\gamma,\beta<\gamma,\delta<\gamma,\alpha_1+\alpha_2+\beta+\delta=2\gamma$
$(1,2;1;1;2)$	$\alpha_1+\alpha_2<\gamma,\alpha_2+\beta<\gamma,\alpha_2+\delta<\gamma,\alpha_1+2\alpha_2+\beta+\delta=2\gamma$

3. Poset $\mathcal{N}=(2;2;1)$



Dimensions D	*Conditions C*
$(0,0;0,0;0;1)$	$\gamma=0$
$(0,1;0,0;0;1)$	$\alpha_2=\gamma$
$(0,0;0,0;1;1)$	$\delta=\gamma$
$(0,1;0,0;1;1)$	$\alpha_2+\delta=\gamma$
$(1,1;0,0;1;1)$	$\alpha_1+\alpha_2+\delta=\gamma$
$(0,0;0,1;0;1)$	$\beta_2=\gamma$
$(0,1;0,1;1;1)$	$\alpha_2+\beta_2+\delta=\gamma$
$(0,0;1,1;0;1)$	$\beta_1+\beta_2=\gamma$
$(0,1;1,1;0;1)$	$\alpha_2+\beta_1+\beta_2=\gamma$
$(0,0;0,1;1;1)$	$\beta_2+\delta=\gamma$
$(0,0;1,1;1;1)$	$\beta_1+\beta_2+\delta=\gamma$
$(0,1;1,1;1;1)$	$\alpha_2+\beta_1+\beta_2+\delta=\gamma$
$(1,1;0,0;0;1)$	$\alpha_1+\alpha_2=\gamma$
$(0,1;0,1;0;1)$	$\alpha_2+\beta_2=\gamma$
$(1,1;0,1;0;1)$	$\alpha_1+\alpha_2+\beta_2=\gamma$
$(1,1;0,1;1;1)$	$\alpha_1+\alpha_2+\beta_2+\delta=\gamma$
$(1,1;1,1;0;1)$	$\alpha_1+\alpha_2+\beta_1+\beta_2=\gamma$
$(1,1;1,1;1;1)$	$\alpha_1+\alpha_2+\beta_1+\beta_2+\delta=\gamma$
$(0,1;0,1;1;2)$	$\alpha_2<\gamma,\beta_2<\gamma,\delta<\gamma,\alpha_2+\beta_2+\delta=2\gamma$
$(1,1;0,1;1;2)$	$\alpha_1+\alpha_2<\gamma,\alpha_1+\alpha_2<\gamma,\beta_2<\gamma,\delta<\gamma,\alpha_1+\alpha_2+\beta_2+\delta=2\gamma$
$(1,2;0,1;1;2)$	$\alpha_1+\alpha_2<\gamma,\alpha_2+\beta_2<\gamma,\alpha_2+\delta<\gamma,\alpha_1+2\alpha_2+\beta_2+\delta=2\gamma$
$(0,1;1,1;1;2)$	$\alpha_2<\gamma,\beta_1+\beta_2<\gamma,\beta_1+\beta_2<\gamma,\delta<\gamma,\alpha_2+\beta_1+\beta_2+\delta=2\gamma$

$(1,1;1,1;1;2)$	$\alpha_1+\alpha_2<\gamma, \alpha_1+\alpha_2<\gamma, \beta_1+\beta_2<\gamma, \beta_1+\beta_2<\gamma, \delta<\gamma, \alpha_1+\alpha_2+\beta_1+\beta_2+\delta=2\gamma$
$(1,2;1,1;1;2)$	$\alpha_1+\alpha_2<\gamma, \alpha_2+\beta_1+\beta_2<\gamma, \alpha_2+\beta_1+\beta_2<\gamma, \alpha_2+\delta<\gamma, \alpha_1+2\alpha_2+\beta_1+\beta_2+\delta=2\gamma$
$(0,1;1,2;1;2)$	$\alpha_2+\beta_2<\gamma, \beta_1+\beta_2<\gamma, \beta_2+\delta<\gamma, \alpha_2+\beta_1+2\beta_2+\delta=2\gamma$
$(1,1;1,2;1;2)$	$\alpha_1+\alpha_2+\beta_2<\gamma, \alpha_1+\alpha_2+\beta_2<\gamma, \beta_1+\beta_2<\gamma, \beta_2+\delta<\gamma, \alpha_1+\alpha_2+\beta_1+2\beta_2+\delta=2\gamma$
$(1,2;1,2;1;2)$	$\alpha_1+\alpha_2+\beta_2<\gamma, \alpha_2+\beta_1+\beta_2<\gamma, \alpha_2+\beta_2+\delta<\gamma, \alpha_1+2\alpha_2+\beta_1+2\beta_2+\delta=2\gamma$
$(1,2;1,2;1;3)$	$\alpha_1+\alpha_2<\gamma, \alpha_1+\alpha_2+\beta_2+\delta<2\gamma, \beta_1+\beta_2<\gamma, \alpha_2+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+\beta_2<\gamma, \alpha_1+2\alpha_2+\beta_1+2\beta_2+\delta=3\gamma$
$(1,2;1,2;2;3)$	$\beta_2+\delta<\gamma, \alpha_2+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+\delta<\gamma, \alpha_1+2\alpha_2+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+\beta_1+2\beta_2+2\delta=3\gamma$

4. Poset $\mathcal{N}=(3;2;1)$



Dimensions D	*Conditions C*
$(0,0,0;0,0;0;1)$	$\gamma=0$
$(0,0,1;0,0;0;1)$	$\alpha_3=\gamma$
$(1,1,1;0,0;0;1)$	$\alpha_1+\alpha_2+\alpha_3=\gamma$
$(0,0,0;0,1;0;1)$	$\beta_2=\gamma$
$(0,0,1;0,1;0;1)$	$\alpha_3+\beta_2=\gamma$
$(0,1,1;0,0;0;1)$	$\alpha_2+\alpha_3=\gamma$
$(0,1,1;0,1;0;1)$	$\alpha_2+\alpha_3+\beta_2=\gamma$
$(1,1,1;0,1;0;1)$	$\alpha_1+\alpha_2+\alpha_3+\beta_2=\gamma$
$(0,0,0;1,1;0;1)$	$\beta_1+\beta_2=\gamma$
$(0,0,1;1,1;0;1)$	$\alpha_3+\beta_1+\beta_2=\gamma$
$(0,1,1;1,1;0;1)$	$\alpha_2+\alpha_3+\beta_1+\beta_2=\gamma$
$(1,1,1;1,1;0;1)$	$\alpha_1+\alpha_2+\alpha_3+\beta_1+\beta_2=\gamma$
$(0,0,0;0,0;1;1)$	$\delta=\gamma$
$(0,0,1;0,0;1;1)$	$\alpha_3+\delta=\gamma$
$(0,0,0;0,1;1;1)$	$\beta_2+\delta=\gamma$
$(0,0,1;0,1;1;1)$	$\alpha_3+\beta_2+\delta=\gamma$
$(0,1,1;0,1;1;1)$	$\alpha_2+\alpha_3+\beta_2+\delta=\gamma$

$(1,1,1;0,1;1;1)$	$\alpha_1+\alpha_2+\alpha_3+\beta_2+\delta=\gamma$
$(0,0,0;1,1;1;1)$	$\beta_1+\beta_2+\delta=\gamma$
$(0,0,1;1,1;1;1)$	$\alpha_3+\beta_1+\beta_2+\delta=\gamma$
$(0,1,1;0,0;1;1)$	$\alpha_2+\alpha_3+\delta=\gamma$
$(0,1,1;1,1;1;1)$	$\alpha_2+\alpha_3+\beta_1+\beta_2+\delta=\gamma$
$(1,1,1;0,0;1;1)$	$\alpha_1+\alpha_2+\alpha_3+\delta=\gamma$
$(1,1,1;1,1;1;1)$	$\alpha_1+\alpha_2+\alpha_3+\beta_1+\beta_2+\delta=\gamma$
$(0,0,1;0,1;1;2)$	$\alpha_3<\gamma,\beta_2<\gamma,\delta<\gamma,\alpha_3+\beta_2+\delta=2\gamma$
$(0,1,1;0,1;1;2)$	$\alpha_2+\alpha_3<\gamma,\alpha_2+\alpha_3<\gamma,\beta_2<\gamma,\delta<\gamma,\alpha_2+\alpha_3+\beta_2+\delta=2\gamma$
$(0,1,2;0,1;1;2)$	$\alpha_2+\alpha_3<\gamma,\alpha_3+\beta_2<\gamma,\alpha_3+\delta<\gamma,\alpha_2+2\alpha_3+\beta_2+\delta=2\gamma$
$(0,0,1;1,1;1;2)$	$\alpha_3<\gamma,\beta_1+\beta_2<\gamma,\beta_1+\beta_2<\gamma,\delta<\gamma,\alpha_3+\beta_1+\beta_2+\delta=2\gamma$
$(0,1,1;1,1;1;2)$	$\alpha_2+\alpha_3<\gamma,\alpha_2+\alpha_3<\gamma,\beta_1+\beta_2<\gamma,\beta_1+\beta_2<\gamma,\delta<\gamma,\alpha_2+\alpha_3+\beta_1+\beta_2+\delta=2\gamma$
$(0,1,2;1,1;1;2)$	$\alpha_2+\alpha_3<\gamma,\alpha_3+\beta_1+\beta_2<\gamma,\alpha_3+\beta_1+\beta_2<\gamma,\alpha_3+\delta<\gamma,\alpha_2+2\alpha_3+\beta_1+\beta_2+\delta=2\gamma$
$(0,1,2;1,2;1;2)$	$\alpha_2+\alpha_3+\beta_2<\gamma,\alpha_3+\beta_1+\beta_2<\gamma,\alpha_3+\beta_2+\delta<\gamma,\alpha_2+2\alpha_3+\beta_1+2\beta_2+\delta=2\gamma$
$(1,1,2;0,1;1;2)$	$\alpha_1+\alpha_2+\alpha_3<\gamma,\alpha_1+\alpha_2+\alpha_3<\gamma,\alpha_3+\beta_2<\gamma,\alpha_3+\delta<\gamma,\alpha_1+\alpha_2+2\alpha_3+\beta_2+\delta=2\gamma$
$(1,1,2;1,1;1;2)$	$\alpha_1+\alpha_2+\alpha_3<\gamma,\alpha_1+\alpha_2+\alpha_3<\gamma,\alpha_3+\beta_1+\beta_2<\gamma,\alpha_3+\beta_1+\beta_2<\gamma,\alpha_3+\delta<\gamma,\alpha_1+\alpha_2+2\alpha_3+\beta_1+\beta_2+\delta=2\gamma$
$(0,0,1;1,2;1;2)$	$\alpha_3+\beta_2<\gamma,\beta_1+\beta_2<\gamma,\beta_2+\delta<\gamma,\alpha_3+\beta_1+2\beta_2+\delta=2\gamma$
$(0,1,1;1,2;1;2)$	$\alpha_2+\alpha_3+\beta_2<\gamma,\alpha_2+\alpha_3+\beta_2<\gamma,\beta_1+\beta_2<\gamma,\beta_2+\delta<\gamma,\alpha_2+\alpha_3+\beta_1+2\beta_2+\delta=2\gamma$
$(1,1,1;0,1;1;2)$	$\alpha_1+\alpha_2+\alpha_3<\gamma,\alpha_1+\alpha_2+\alpha_3<\gamma,\alpha_1+\alpha_2+\alpha_3<\gamma,\beta_2<\gamma,\delta<\gamma,\alpha_1+\alpha_2+\alpha_3+\beta_2+\delta=2\gamma$
$(1,1,1;1,1;1;2)$	$\alpha_1+\alpha_2+\alpha_3<\gamma,\alpha_1+\alpha_2+\alpha_3<\gamma,\alpha_1+\alpha_2+\alpha_3<\gamma,\beta_1+\beta_2<\gamma,\beta_1+\beta_2<\gamma,\delta<\gamma,\alpha_1+\alpha_2+\alpha_3+\beta_1+\beta_2+\delta=2\gamma$
$(1,1,1;1,2;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\beta_2<\gamma,\alpha_1+\alpha_2+\alpha_3+\beta_2<\gamma,\alpha_1+\alpha_2+\alpha_3+\beta_2<\gamma,\beta_1+\beta_2<\gamma,\beta_2+\delta<\gamma,\alpha_1+\alpha_2+\alpha_3+\beta_1+2\beta_2+\delta=2\gamma$
$(1,1,2;1,2;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\beta_2<\gamma,\alpha_1+\alpha_2+\alpha_3+\beta_2<\gamma,\alpha_3+\beta_1+\beta_2<\gamma,\alpha_3+\beta_2+\delta<\gamma,\alpha_1+\alpha_2+2\alpha_3+\beta_1+2\beta_2+\delta=2\gamma$

$(1,2,2;0,1;1;2)$	$\alpha_1+\alpha_2+\alpha_3<\gamma, \alpha_2+\alpha_3+\beta_2<\gamma, \alpha_2+\alpha_3+\delta<\gamma, \alpha_1+2\alpha_2+2\alpha_3+\beta_2+\delta=2\gamma$
$(1,2,2;1,1;1;2)$	$\alpha_1+\alpha_2+\alpha_3<\gamma, \alpha_2+\alpha_3+\beta_1+\beta_2<\gamma, \alpha_2+\alpha_3+\beta_1+\beta_2<\gamma, \alpha_2+\alpha_3+\delta<\gamma, \alpha_1+2\alpha_2+2\alpha_3+\beta_1+\beta_2+\delta=2\gamma$
$(1,2,2;1,2;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\beta_2<\gamma, \alpha_2+\alpha_3+\beta_1+\beta_2<\gamma, \alpha_2+\alpha_3+\beta_2+\delta<\gamma, \alpha_1+2\alpha_2+2\alpha_3+\beta_1+2\beta_2+\delta=2\gamma$
$(0,1,2;1,2;1;3)$	$\alpha_2+\alpha_3<\gamma, \alpha_2+\alpha_3+\beta_2+\delta<2\gamma, \beta_1+\beta_2<\gamma, \alpha_3+\beta_1+\beta_2+\delta<2\gamma, \alpha_3+\beta_2<\gamma, \alpha_2+2\alpha_3+\beta_1+2\beta_2+\delta=3\gamma$
$(1,1,2;1,2;1;3)$	$\alpha_1+\alpha_2+\alpha_3<\gamma, \alpha_1+\alpha_2+\alpha_3<\gamma, \alpha_1+\alpha_2+\alpha_3+\beta_2+\delta<2\gamma, \beta_1+\beta_2<\gamma, \alpha_3+\beta_1+\beta_2+\delta<2\gamma, \alpha_3+\beta_2<\gamma, \alpha_1+\alpha_2+2\alpha_3+\beta_1+2\beta_2+\delta=3\gamma$
$(1,2,2;1,2;1;3)$	$\alpha_1+\alpha_2+\alpha_3<\gamma, \alpha_1+\alpha_2+\alpha_3+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+\beta_2+\delta<2\gamma, \beta_1+\beta_2<\gamma, \alpha_2+\alpha_3+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\beta_2<\gamma, \alpha_1+2\alpha_2+2\alpha_3+\beta_1+2\beta_2+\delta=3\gamma$
$(1,2,3;1,2;1;3)$	$\alpha_1+\alpha_2+\alpha_3<\gamma, \alpha_1+\alpha_2+2\alpha_3+\beta_2+\delta<2\gamma, \alpha_3+\beta_1+\beta_2<\gamma, \alpha_2+2\alpha_3+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\beta_2<\gamma, \alpha_1+2\alpha_2+3\alpha_3+\beta_1+2\beta_2+\delta=3\gamma$
$(0,1,2;1,2;2;3)$	$\beta_2+\delta<\gamma, \alpha_3+\beta_1+2\beta_2+\delta<2\gamma, \alpha_3+\delta<\gamma, \alpha_2+2\alpha_3+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+2\alpha_3+\beta_1+2\beta_2+2\delta=3\gamma$
$(1,1,2;1,2;2;3)$	$\beta_2+\delta<\gamma, \beta_2+\delta<\gamma, \alpha_3+\beta_1+2\beta_2+\delta<2\gamma, \alpha_3+\delta<\gamma, \alpha_1+\alpha_2+2\alpha_3+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+2\alpha_3+\beta_1+2\beta_2+2\delta=3\gamma$
$(1,2,2;1,2;2;3)$	$\beta_2+\delta<\gamma, \alpha_2+\alpha_3+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\delta<\gamma, \alpha_1+2\alpha_2+2\alpha_3+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+2\alpha_3+\beta_1+2\beta_2+2\delta=3\gamma$
$(1,2,3;1,2;2;3)$	$\alpha_3+\beta_2+\delta<\gamma, \alpha_2+2\alpha_3+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\delta<\gamma, \alpha_1+2\alpha_2+2\alpha_3+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+2\alpha_3+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+3\alpha_3+\beta_1+2\beta_2+2\delta=3\gamma$
$(1,2,3;1,2;2;4)$	$\alpha_1+\alpha_2+\alpha_3<\gamma, \alpha_1+\alpha_2+\alpha_3+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+2\alpha_3+\beta_1+2\beta_2+\delta<3\gamma, \alpha_3+\delta<\gamma, \alpha_2+2\alpha_3+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+3\alpha_3+\beta_1+2\beta_2+2\delta=4\gamma$

$(1,2,3;1,3;2;4)$	$\beta_2+\delta<\gamma, \alpha_3+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+2\alpha_3+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_2+\alpha_3+\beta_2<\gamma, \alpha_1+2\alpha_2+2\alpha_3+\beta_1+2\beta_2+\delta<3\gamma, \alpha_1+\alpha_2+2\alpha_3+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+3\alpha_3+\beta_1+3\beta_2+2\delta=4\gamma$
$(1,2,3;2,3;2;4)$	$\alpha_3+\beta_1+\beta_2<\gamma, \alpha_2+2\alpha_3+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+3\alpha_3+\beta_1+2\beta_2+\delta<3\gamma, \alpha_1+\alpha_2+\alpha_3+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+2\alpha_3+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_2+\alpha_3+\beta_1+2\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+3\alpha_3+2\beta_1+3\beta_2+2\delta=4\gamma$

5. Poset $\mathcal{N}=(4;2;1)$



Dimensions D	*Conditions C*
$(0,0,0,0;0,0;0;1)$	$\gamma=0$
$(1,1,1,1;0,0;1;1)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4+\delta=\gamma$
$(0,0,0,0;0,1;0;1)$	$\beta_2=\gamma$
$(0,0,0,0;1,1;1;1)$	$\beta_1+\beta_2+\delta=\gamma$
$(0,0,0,1;0,0;0;1)$	$\alpha_4=\gamma$
$(0,0,0,0;0,0;1;1)$	$\delta=\gamma$
$(0,0,0,1;0,0;1;1)$	$\alpha_4+\delta=\gamma$
$(0,0,0,0;0,1;1;1)$	$\beta_2+\delta=\gamma$
$(0,0,0,1;0,1;0;1)$	$\alpha_4+\beta_2=\gamma$
$(0,0,0,1;0,1;1;1)$	$\alpha_4+\beta_2+\delta=\gamma$
$(0,0,0,0;1,1;0;1)$	$\beta_1+\beta_2=\gamma$
$(0,0,0,1;1,1;0;1)$	$\alpha_4+\beta_1+\beta_2=\gamma$
$(0,0,0,1;1,1;1;1)$	$\alpha_4+\beta_1+\beta_2+\delta=\gamma$
$(0,0,1,1;0,0;0;1)$	$\alpha_3+\alpha_4=\gamma$
$(0,0,1,1;0,0;1;1)$	$\alpha_3+\alpha_4+\delta=\gamma$
$(0,0,1,1;1,1;1;1)$	$\alpha_3+\alpha_4+\beta_1+\beta_2+\delta=\gamma$
$(0,1,1,1;0,0;1;1)$	$\alpha_2+\alpha_3+\alpha_4+\delta=\gamma$
$(0,0,1,1;0,1;1;1)$	$\alpha_3+\alpha_4+\beta_2+\delta=\gamma$
$(0,1,1,1;0,1;1;1)$	$\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta=\gamma$
$(0,1,1,1;1,1;1;1)$	$\alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta=\gamma$
$(1,1,1,1;0,1;1;1)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta=\gamma$
$(0,0,1,1;1,1;0;1)$	$\alpha_3+\alpha_4+\beta_1+\beta_2=\gamma$
$(0,1,1,1;0,0;0;1)$	$\alpha_2+\alpha_3+\alpha_4=\gamma$
$(0,0,1,1;0,1;0;1)$	$\alpha_3+\alpha_4+\beta_2=\gamma$

$(0,1,1,1;0,1;0;1)$	$\alpha_2+\alpha_3+\alpha_4+\beta_2=\gamma$
$(0,1,1,1;1,1;0;1)$	$\alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2=\gamma$
$(1,1,1,1;0,0;0;1)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4=\gamma$
$(1,1,1,1;0,1;0;1)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2=\gamma$
$(1,1,1,1;1,1;0;1)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2=\gamma$
$(1,1,1,1;1,1;1;1)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta=\gamma$
$(0,0,0,1;0,1;1;2)$	$\alpha_4<\gamma,\beta_2<\gamma,\delta<\gamma,\alpha_4+\beta_2+\delta=2\gamma$
$(0,0,1,1;0,1;1;2)$	$\alpha_3+\alpha_4<\gamma,\alpha_3+\alpha_4<\gamma,\beta_2<\gamma,\delta<\gamma,\alpha_3+\alpha_4+\beta_2+\delta=2\gamma$
$(1,1,1,1;0,1;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma,\beta_2<\gamma,\delta<\gamma,\alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta=2\gamma$
$(0,0,1,2;0,1;1;2)$	$\alpha_3+\alpha_4<\gamma,\alpha_4+\beta_2<\gamma,\alpha_4+\delta<\gamma,\alpha_3+2\alpha_4+\beta_2+\delta=2\gamma$
$(0,1,1,1;0,1;1;2)$	$\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_2+\alpha_3+\alpha_4<\gamma,\beta_2<\gamma,\delta<\gamma,\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta=2\gamma$
$(1,1,1,2;0,1;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_4+\beta_2<\gamma,\alpha_4+\delta<\gamma,\alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta=2\gamma$
$(1,1,2,2;0,1;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_3+\alpha_4+\beta_2<\gamma,\alpha_3+\alpha_4+\delta<\gamma,\alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta=2\gamma$
$(1,2,2,2;0,1;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma,\alpha_2+\alpha_3+\alpha_4+\delta<\gamma,\alpha_1+2\alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta=2\gamma$
$(0,0,0,1;1,1;1;2)$	$\alpha_4<\gamma,\beta_1+\beta_2<\gamma,\beta_1+\beta_2<\gamma,\delta<\gamma,\alpha_4+\beta_1+\beta_2+\delta=2\gamma$
$(0,0,1,1;1,1;1;2)$	$\alpha_3+\alpha_4<\gamma,\alpha_3+\alpha_4<\gamma,\beta_1+\beta_2<\gamma,\beta_1+\beta_2<\gamma,\delta<\gamma,\alpha_3+\alpha_4+\beta_1+\beta_2+\delta=2\gamma$
$(0,0,1,2;1,1;1;2)$	$\alpha_3+\alpha_4<\gamma,\alpha_4+\beta_1+\beta_2<\gamma,\alpha_4+\beta_1+\beta_2<\gamma,\alpha_4+\delta<\gamma,\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta=2\gamma$
$(0,1,1,1;1,1;1;2)$	$\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_2+\alpha_3+\alpha_4<\gamma,\beta_1+\beta_2<\gamma,\beta_1+\beta_2<\gamma,\delta<\gamma,\alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta=2\gamma$
$(1,1,1,1;1,1;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma,\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma,\beta_1+\beta_2<\gamma,\beta_1+\beta_2<\gamma,\delta<\gamma,\alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta=2\gamma$

$(1,1,1,2;1,1;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_4+\beta_1+\beta_2<\gamma, \alpha_4+\beta_1+\beta_2<\gamma, \alpha_4+\delta<\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta=2\gamma$
$(0,1,2,2;0,1;1;2)$	$\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_3+\alpha_4+\beta_2<\gamma, \alpha_3+\alpha_4+\delta<\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta=2\gamma$
$(0,1,2,2;1,1;1;2)$	$\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_3+\alpha_4+\beta_1+\beta_2<\gamma, \alpha_3+\alpha_4+\beta_1+\beta_2<\gamma, \alpha_3+\alpha_4+\delta<\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta=2\gamma$
$(1,1,2,2;1,1;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_3+\alpha_4+\beta_1+\beta_2<\gamma, \alpha_3+\alpha_4+\beta_1+\beta_2<\gamma, \alpha_3+\alpha_4+\delta<\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta=2\gamma$
$(1,2,2,2;1,1;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2<\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2<\gamma, \alpha_2+\alpha_3+\alpha_4+\delta<\gamma, \alpha_1+2\alpha_2+2\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta=2\gamma$
$(0,0,0,1;1,2;1;2)$	$\alpha_4+\beta_2<\gamma, \beta_1+\beta_2<\gamma, \beta_2+\delta<\gamma, \alpha_4+\beta_1+2\beta_2+\delta=2\gamma$
$(0,0,1,1;1,2;1;2)$	$\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_3+\alpha_4+\beta_2<\gamma, \beta_1+\beta_2<\gamma, \beta_2+\delta<\gamma, \alpha_3+\alpha_4+\beta_1+2\beta_2+\delta=2\gamma$
$(0,0,1,2;1,2;1;2)$	$\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_4+\beta_1+\beta_2<\gamma, \alpha_4+\beta_2+\delta<\gamma, \alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta=2\gamma$
$(0,1,1,2;0,1;1;2)$	$\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_4+\beta_2<\gamma, \alpha_4+\delta<\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta=2\gamma$
$(0,1,1,2;1,1;1;2)$	$\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_4+\beta_1+\beta_2<\gamma, \alpha_4+\beta_1+\beta_2<\gamma, \alpha_4+\delta<\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta=2\gamma$
$(0,1,1,1;1,2;1;2)$	$\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \beta_1+\beta_2<\gamma, \beta_2+\delta<\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+2\beta_2+\delta=2\gamma$
$(0,1,1,2;1,2;1;2)$	$\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_4+\beta_1+\beta_2<\gamma, \alpha_4+\beta_2+\delta<\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta=2\gamma$
$(1,1,1,2;1,2;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_4+\beta_1+\beta_2<\gamma, \alpha_4+\beta_2+\delta<\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta=2\gamma$
$(0,1,2,2;1,2;1;2)$	$\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_3+\alpha_4+\beta_1+\beta_2<\gamma, \alpha_3+\alpha_4+\beta_2+\delta<\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta=2\gamma$

$(1,1,1,1;1,2;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \beta_1+\beta_2<\gamma, \beta_2+\delta<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_1+2\beta_2+\delta=2\gamma$
$(1,1,2,2;1,2;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_3+\alpha_4+\beta_1+\beta_2<\gamma, \alpha_3+\alpha_4+\beta_2+\delta<\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta=2\gamma$
$(1,2,2,2;1,2;1;2)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2<\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<\gamma, \alpha_1+2\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta=2\gamma$
$(0,0,1,2;1,2;1;3)$	$\alpha_3+\alpha_4<\gamma, \alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \beta_1+\beta_2<\gamma, \alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_4+\beta_2<\gamma, \alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta=3\gamma$
$(0,1,1,2;1,2;1;3)$	$\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \beta_1+\beta_2<\gamma, \alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_4+\beta_2<\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta=3\gamma$
$(0,1,2,2;1,2;1;3)$	$\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \beta_1+\beta_2<\gamma, \alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_3+\alpha_4+\beta_2<\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta=3\gamma$
$(0,0,1,2;1,2;2;3)$	$\beta_2+\delta<\gamma, \alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_4+\delta<\gamma, \alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta=3\gamma$
$(0,1,1,2;1,2;2;3)$	$\beta_2+\delta<\gamma, \beta_2+\delta<\gamma, \alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_4+\delta<\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta=3\gamma$
$(0,1,2,2;1,2;2;3)$	$\beta_2+\delta<\gamma, \alpha_3+\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_3+\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_3+\alpha_4+\delta<\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta=3\gamma$
$(0,1,2,3;1,2;1;3)$	$\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_4+\beta_1+\beta_2<\gamma, \alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_3+\alpha_4+\beta_2<\gamma, \alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta=3\gamma$
$(0,1,2,3;1,2;2;3)$	$\alpha_4+\beta_2+\delta<\gamma, \alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_3+\alpha_4+\delta<\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta=3\gamma$

$(1,1,1,2;1,2;1;3)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \beta_1+\beta_2<\gamma, \alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_4+\beta_2<\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta=3\gamma$
$(1,1,2,2;1,2;1;3)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \beta_1+\beta_2<\gamma, \alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta=3\gamma$
$(1,1,2,3;1,2;1;3)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_4+\beta_1+\beta_2<\gamma, \alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta=3\gamma$
$(1,1,1,2;1,2;2;3)$	$\beta_2+\delta<\gamma, \beta_2+\delta<\gamma, \beta_2+\delta<\gamma, \alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_4+\delta<\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta=3\gamma$
$(1,1,2,2;1,2;2;3)$	$\beta_2+\delta<\gamma, \beta_2+\delta<\gamma, \alpha_3+\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_3+\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_3+\alpha_4+\delta<\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta=3\gamma$
$(1,1,2,3;1,2;2;3)$	$\alpha_4+\beta_2+\delta<\gamma, \alpha_4+\beta_2+\delta<\gamma, \alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_3+\alpha_4+\delta<\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta=3\gamma$
$(1,2,2,2;1,2;1;3)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \beta_1+\beta_2<\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+2\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta=3\gamma$
$(1,2,2,3;1,2;1;3)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_4+\beta_1+\beta_2<\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta=3\gamma$
$(1,2,2,2;1,2;2;3)$	$\beta_2+\delta<\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\alpha_4+\delta<\gamma, \alpha_1+2\alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta=3\gamma$

$(1,2,2,3;1,2;2;3)$	$\alpha_4+\beta_2+\delta<\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\alpha_4+\delta<\gamma, \alpha_1+2\alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta=3\gamma$
$(1,2,3,3;1,2;1;3)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_3+\alpha_4+\beta_1+\beta_2<\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+2\alpha_2+3\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta=3\gamma$
$(1,2,3,3;1,2;2;3)$	$\alpha_3+\alpha_4+\beta_2+\delta<\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\alpha_4+\delta<\gamma, \alpha_1+2\alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+3\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta=3\gamma$
$(0,1,2,3;1,2;2;4)$	$\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_4+\delta<\gamma, \alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta=4\gamma$
$(1,1,2,3;1,2;2;4)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_4+\delta<\gamma, \alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta=4\gamma$
$(1,2,2,3;1,2;2;4)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_4+\delta<\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta=4\gamma$
$(1,2,3,3;1,2;2;4)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_3+\alpha_4+\delta<\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+3\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta=4\gamma$
$(1,2,3,4;1,2;2;4)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4<\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_3+\alpha_4+\delta<\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+\beta_1+2\beta_2+2\delta=4\gamma$
$(0,1,2,3;1,3;2;4)$	$\beta_2+\delta<\gamma, \alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_3+\alpha_4+\beta_2<\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_2+2\alpha_3+3\alpha_4+\beta_1+3\beta_2+2\delta=4\gamma$

$(1,1,2,3;1,3;2;4)$	$\beta_2+\delta<\gamma, \beta_2+\delta<\gamma, \alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+\beta_1+3\beta_2+2\delta=4\gamma$
$(1,2,2,3;1,3;2;4)$	$\beta_2+\delta<\gamma, \alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+2\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+\beta_1+3\beta_2+2\delta=4\gamma$
$(1,2,3,3;1,3;2;4)$	$\beta_2+\delta<\gamma, \alpha_3+\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+2\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+3\alpha_3+3\alpha_4+\beta_1+3\beta_2+2\delta=4\gamma$
$(1,2,3,4;1,3;2;4)$	$\alpha_4+\beta_2+\delta<\gamma, \alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+\beta_1+3\beta_2+2\delta=4\gamma$
$(0,1,2,3;2,3;2;4)$	$\alpha_4+\beta_1+\beta_2<\gamma, \alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_3+\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+2\alpha_3+3\alpha_4+2\beta_1+3\beta_2+2\delta=4\gamma$
$(1,1,2,3;2,3;2;4)$	$\alpha_4+\beta_1+\beta_2<\gamma, \alpha_4+\beta_1+\beta_2<\gamma, \alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_3+\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+2\beta_1+3\beta_2+2\delta=4\gamma$
$(1,2,2,3;2,3;2;4)$	$\alpha_4+\beta_1+\beta_2<\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+2\beta_1+3\beta_2+2\delta=4\gamma$

$(1,2,3,3;2,3;2;4)$	$\alpha_3+\alpha_4+\beta_1+\beta_2 < \gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta < 2\gamma, \alpha_1+2\alpha_2+3\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta < 3\gamma, \alpha_1+2\alpha_2+3\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta < 3\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta < 2\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta < 3\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+2\beta_2+\delta < 2\gamma, \alpha_1+2\alpha_2+3\alpha_3+3\alpha_4+2\beta_1+3\beta_2+2\delta = 4\gamma$
$(1,2,3,4;2,3;2;4)$	$\alpha_3+\alpha_4+\beta_1+\beta_2 < \gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta < 2\gamma, \alpha_1+2\alpha_2+3\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta < 3\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta < 2\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta < 3\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta < 2\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+2\beta_1+3\beta_2+2\delta = 4\gamma$
$(1,2,3,4;1,3;2;5)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4 < \gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_2+\delta < 2\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta < 3\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta < 4\gamma, \alpha_3+\alpha_4+\beta_2 < \gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta < 3\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta < 2\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+\beta_1+3\beta_2+2\delta = 5\gamma$
$(1,2,3,4;2,3;2;5)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4 < \gamma, \alpha_4+\beta_1+\beta_2 < \gamma, \alpha_3+2\alpha_4+\beta_1+\beta_2+\delta < 2\gamma, \alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta < 3\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta < 2\gamma, \alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta < 4\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta < 3\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+2\beta_1+3\beta_2+2\delta = 5\gamma$
$(1,2,3,4;2,4;2;5)$	$\alpha_4+\beta_1+\beta_2 < \gamma, \alpha_2+\alpha_3+\alpha_4+\beta_2 < \gamma, \alpha_1+2\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta < 3\gamma, \alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+\beta_1+3\beta_2+2\delta < 4\gamma, \alpha_3+\alpha_4+\beta_1+2\beta_2+\delta < 2\gamma, \alpha_2+2\alpha_3+3\alpha_4+2\beta_1+3\beta_2+2\delta < 4\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta < 3\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+2\beta_1+4\beta_2+2\delta = 5\gamma$
$(1,2,3,4;1,3;3;5)$	$\beta_2+\delta < \gamma, \alpha_3+\alpha_4+\delta < \gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta < 2\gamma, \alpha_1+2\alpha_2+3\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta < 4\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta < 2\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+\beta_1+3\beta_2+2\delta < 4\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta < 3\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+\beta_1+3\beta_2+3\delta = 5\gamma$

$(1,2,3,4;2,3;3;5)$	$\alpha_3+\alpha_4+\delta<\gamma, \alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+2\beta_1+3\beta_2+2\delta<4\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+\beta_1+2\beta_2+2\delta<4\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+2\beta_1+3\beta_2+3\delta=5\gamma$
$(1,2,3,4;2,4;3;5)$	$\alpha_4+\beta_2+\delta<\gamma, \alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+\beta_1+3\beta_2+2\delta<4\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+2\beta_1+3\beta_2+2\delta<4\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+2\beta_1+4\beta_2+3\delta=5\gamma$
$(1,2,3,4;2,4;3;6)$	$\beta_2+\delta<\gamma, \alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_2+2\alpha_3+3\alpha_4+\beta_1+3\beta_2+2\delta<4\gamma, \alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta<4\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+2\beta_1+4\beta_2+3\delta=6\gamma$
$(1,2,3,5;2,4;3;6)$	$\alpha_4+\beta_1+\beta_2<\gamma, \alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+2\beta_1+3\beta_2+2\delta<5\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+\beta_1+3\beta_2+2\delta<4\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_1+2\alpha_2+3\alpha_3+5\alpha_4+2\beta_1+4\beta_2+3\delta=6\gamma$
$(1,2,4,5;2,4;3;6)$	$\alpha_3+\alpha_4+\delta<\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+3\alpha_3+3\alpha_4+\beta_1+2\beta_2+2\delta<4\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+\beta_1+3\beta_2+3\delta<5\gamma, \alpha_3+\alpha_4+\beta_1+2\beta_2+\delta<2\gamma, \alpha_2+2\alpha_3+3\alpha_4+2\beta_1+3\beta_2+2\delta<4\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_1+2\alpha_2+4\alpha_3+5\alpha_4+2\beta_1+4\beta_2+3\delta=6\gamma$
$(1,3,4,5;2,4;3;6)$	$\alpha_2+\alpha_3+\alpha_4+\beta_2<\gamma, \alpha_1+2\alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, \alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+\beta_1+3\beta_2+2\delta<4\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+2\beta_1+4\beta_2+2\delta<5\gamma, \alpha_2+\alpha_3+2\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+\beta_1+2\beta_2+2\delta<4\gamma, \alpha_2+2\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_1+3\alpha_2+4\alpha_3+5\alpha_4+2\beta_1+4\beta_2+3\delta=6\gamma$

$(2,3,4,5;2,4;3;6)$	$\alpha_1+\alpha_2+\alpha_3+\alpha_4+\beta_1+\beta_2+\delta<2\gamma, \alpha_1+\alpha_2+\alpha_3+2\alpha_4+\beta_1+2\beta_2+2\delta<3\gamma, \alpha_1+\alpha_2+2\alpha_3+3\alpha_4+2\beta_1+3\beta_2+2\delta<4\gamma, \alpha_1+2\alpha_2+3\alpha_3+4\alpha_4+2\beta_1+3\beta_2+3\delta<5\gamma, \alpha_1+\alpha_2+2\alpha_3+2\alpha_4+\beta_2+\delta<2\gamma, \alpha_1+2\alpha_2+3\alpha_3+3\alpha_4+\beta_1+3\beta_2+2\delta<4\gamma, \alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+\beta_1+2\beta_2+\delta<3\gamma, 2\alpha_1+3\alpha_2+4\alpha_3+5\alpha_4+2\beta_1+4\beta_2+3\delta=6\gamma$

Acknowledgments. The authors are deeply appreciate to Prof. Yu.Samoilenko for the statement of problem and for his constant attention to this work.

References

[1] Nazarova L. A., Roiter A. V. *Representations of the partially ordered sets,*Investigations on the theory of representations. Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)(Russian) **28** (1972), P.5–31.

[2] Kleiner M. M. *Partially ordered sets of finite type,* Investigations on the theory of representations. Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)(Russian) **28** (1972), P.32–41.

[3] Kleiner M. M. *On the faithful represenattions of partially ordered sets of finite type,* Investigations on the theory of representations. Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)(Russian) **28** (1972), P.42–59.

[4] Drozd Ju. A. *Coxeter transformations and representations of partially ordered sets,* Funkcional. Anal. i Prilozen. (Russian) **8** (1974), no. 3, P.34–42.

[5] Krugljak S. A., Samoilenko Yu. S. *Unitary equivalence of sets of selfajoint operators,* Funkcional. Anal. i Prilozen. (Russian) **14**, (1980), no. 1, P.60–62.

[6] Krugljak S. A., Samoilenko Yu. S. *On the complexity of description of representations of $*$-algebras generated by idempotents,* Proceedings of the American Mathematical Society **128**, (2000), no. 6, P.1655–1664.

[7] Gabriel P., Roiter A. V. *Representations of Finite-Dimensional Algebras,* Enc. of Math. Sci., **73**, Algebra VIII, (Springer, 1992).

[8] Assem I., Simpson D., Skowronskiy A. *Elements of the Representation Theory of Assosiative Algebras. Volume 1 Techniques of Representation Theory,* London Mathematical Society Student Texts 65, (2006), P.458.

[9] Dlab V. and Ringel C.M. *Indecomposable representations of graphs and algebras,* Memoris Amer. Math. Soc, **6**, (1976), no. 176, P.1–72.

[10] P. Donovan and M. Freislich *The Representation Theory of Finite Graphs and Associated Algebras,* Carleton Mathematical Lecture Notes, 1973, **5**, P.1–187.

[11] M. Enomoto and Ya. Watatani *Relative position of four subspaces in a Hilbert space,* Adv. Math., **201**, (2006), no. 2, P.263–317.

[12] Berstein I.N., Gelfand I. M., Ponomarev V. A. *Coxeter functors, and Gabriel's theorem,* Uspehi Mat. Nauk (Russian) **2**, (1973), no. 170, P.19–33.

[13] S. A. Kruglyak, L. A. Nazarova, A. V. Roiter *Orthoscalar representations of quivers in the category of Hilbert spaces,* J. Math. Sci. (N. Y.), **145**, (2007), no. 1, P.4793–4804.

[14] Kruglyak S. A., Roiter A. V. *Locally scalar representations of graphs in the category of Hilbert spaces,* Funct. Anal. Appl., **39**, (2005), no. 2. P.91–105.

[15] Krugljak S. A., Popovych S. V., Samoilenko Yu. S. *The spectral problem and $*$-representations of algebras associated with Dynkin graphs*, J. Algebra Appl. **4** (2005), no. 6, P.761–776.

[16] Crawley-Boevey W. and Geiss Ch. *Horn's problem and semi-stabilty for quiver representations,* Representations of Algebras, V.I.Poc. IX Intern. Conf. Benjing, (2000), P.40–48.

[17] Ostrovskyi V. L.,Samoilenko Yu. S. *On spectral theorems for families of linearly connected selfadjoint operators with prescribed spectra associated with extended Dynkin graphs* Ukrain. Mat. Zh. (Ukrainian) **58** (2006), no. 11, P.1556–1570.

Roman Grushevoy
Institute of Mathematics NAS of Ukraine
Tereschenkivska str. 3
01601 Kiev
Ukraine
e-mail: `grushevoy@imath.kiev.ua`

Kostyantyn Yusenko
Institute of Mathematics NAS of Ukraine
Tereschenkivska str. 3
01601 Kiev
Ukraine
e-mail: `kay@imath.kiev.ua`